









# New records and geographic distribution map of *Proceratophrys sanctaritae* Cruz & Napoli, 2010 (Anura, Odontophrynidae)

Ana Paula Almeida<sup>1\*</sup>, Deivson Fraga de Oliveira Bastos<sup>1</sup>, Paulo Barros de Abreu Júnior<sup>1</sup>, Marcos Ferreira Vila Nova<sup>2</sup>, Iuri Ribeiro Dias<sup>2</sup>, Juliana Zina<sup>1</sup>

**1** Herpetology Laboratory, Department of Biological Sciences, Universidade Estadual do Sudoeste da Bahia, Jequié, BA, Brazil • APA: [sousa.apa.biologia@gmail.com](mailto:sousa.apa.biologia@gmail.com)  <https://orcid.org/0000-0002-5252-6210> • DFOB: [deivsonfraga@gmail.com](mailto:deivsonfraga@gmail.com)  <https://orcid.org/0000-0003-1333-4432> • PBAJr: [pbarrosbio@gmail.com](mailto:pbarrosbio@gmail.com)  <https://orcid.org/0000-0001-9839-0796> • JZ: [juzina74@gmail.com](mailto:juzina74@gmail.com)  <https://orcid.org/0000-0002-9759-1972>

**2** Tropical Herpetology Lab, Department of Biological Sciences, Universidade Estadual de Santa Cruz, Ilhéus, BA, Brazil • MFVN: [marcos.ferreira.vilanova@gmail.com](mailto:marcos.ferreira.vilanova@gmail.com)  <https://orcid.org/0000-0003-1925-4695> • IRD: [iurirdias@hotmail.com](mailto:iurirdias@hotmail.com)  <https://orcid.org/0000-0002-2825-3494>

\* Corresponding author

## Abstract

Until now, *Proceratophrys sanctaritae* Cruz & Napoli, 2010 has only been known from its type locality, Serra do Timbó, municipality of Amargosa, Bahia, Brazil. Due to its restricted geographic range, this species is considered Critically Endangered. We report new records of *P. sanctaritae* in Bahia and update its geographic distribution. Although the geographic range of *P. sanctaritae* is significantly expanded with our new data, the mountainous forest habitat of this species is highly susceptible to human modification, and *P. sanctaritae* is still likely threatened under IUCN criteria.

## Keywords

Atlantic Forest, Caatinga, conservation, horned frogs, transitional areas

**Academic editor:** Raúl Maneyro | Received 23 March 2022 | Accepted 4 August 2022 | Published 16 November 2022

**Citation:** Almeida AP, Bastos DFO, Abreu Júnior PB, Nova MFV, Dias IR, Zina J (2022) New records and geographic distribution map of *Proceratophrys sanctaritae* Cruz & Napoli, 2010 (Anura, Odontophrynidae). Check List 18 (6): 1243–1247. <https://doi.org/10.15560/18.6.1243>

## Introduction

*Proceratophrys* Miranda-Ribeiro, 1920 is the most species-rich genus of the family Odontophrynidae, with 43 species distributed throughout the Neotropics (Frost 2022). Popularly known as horned frogs, these organisms are found in leaf litter near lotic environments, where their cryptic color pattern makes it difficult to distinguish them (Prado and Pombal 2008).

The taxonomy of *Proceratophrys* is mostly based on morphology, with eyelid and rostral appendages essential in defining species groups in the genus. For example,

the *P. appendiculata* group, to which *P. sanctaritae* Cruz & Napoli, 2010 belongs, is characterized by eyelid appendages and a triangular rostral appendix (Prado and Pombal 2008). Nevertheless, a recent analysis does not support the monophyly proposed for most of the species in the group (Dias et al. 2013). Additional phylogenetic studies, including a larger data set, is necessary to elucidate the evolutionary relationships among *Proceratophrys* species (Santana et al. 2021).

*Proceratophrys sanctaritae* (Fig. 1) is a medium-sized





**Figure 1.** *Proceratophrys sanctaritae* from Amargosa municipality, Bahia, Brazil.

frog (mean male snout–vent length (SVL) = 41.0 mm) that is endemic to the Atlantic Forest in Brazil (Cruz and Napoli 2010). Until 2018, it was known only from its type locality, Floresta Santa Rita (13°04'S, 039°38'W) in the Serra do Timbó, Amargosa municipality, Bahia (Frost 2022). The Serra do Timbó is a mountain range characterized by a rugged relief, between 600 and 900 m a.s.l., and covered mainly by semideciduous tropical forest fragments (Cruz and Napoli 2010). Despite the humid tropical climate, the Serra do Timbó is in a transition zone between the Caatinga and Atlantic Forest morpho-climatic domains (Olson et al. 2001).

*Proceratophrys sanctaritae* was categorized as Critically Endangered, as it was only known from a single location under threat from habitat loss and degradation due to livestock and agriculture. Here, we provide new occurrence records that expand the species' geographic distribution to an ecotone region in south-central Bahia and elevational range.

## Methods

Our new records were collected as part of anurofauna inventories conducted between 2017 and 2019 at two sites in Bahia: Morro do Mara and Wenceslau Guimarães State Ecological Station. Morro do Mara (MM; altitude = 560–960 m) is a Seasonal Semideciduous Forest fragment in good preservation, with lentic and lotic water bodies, and surrounded by cattle pastures and cocoa plantations. The Wenceslau Guimarães State Ecological Station (WGSES; altitude = 440–900 m) is characterized by a Submountain Dense Ombrophilous Forest comprising well-preserved forested fragments surrounded by cattle pastures and cocoa plantations. WGSES is in the Central Atlantic Forest Corridor, an important conservation region with many endemic species and high biodiversity. At both sites, lakes, temporary and permanent ponds, streams, and the litter of trails near these water

bodies were monitored.

Serra do Timbó, WGSES, and MM are hilltops in the transition between the Atlantic Forest and Caatinga, and the three sites are exposed to threats associated with human activities as agriculture expansion, pasturing livestock, and flooding behind dams. We collected our specimens of *P. sanctaritae* under SISBIO collection licenses #35068 and #18.660. Specimens were euthanized with an overdose of 5% lidocaine, fixed in 10% formalin, and stored in 70% alcohol. The specimens were deposited in the zoological collection of the Universidade Estadual do Sudoeste da Bahia (MHNJCH; Jequié, Bahia Brazil) and Zoological Museum of Universidade Estadual de Santa Cruz (MZUESC; Ilhéus, Bahia, Brazil).

The altimetry distribution map was built from elevation data available in the SRTM 90m DEM Digital Elevation Database on the CGIAR-CSI portal (<https://srtm.csi.cgiar.org>). The data present a 90 m resolution at the equator and is available in 5 × 5-degree grids. Data were obtained using the *raster* package v. 3.4.13 (Hijman 2021) and the map produced with packages *rgdal* v. 1.5.21 (Bivand et al. 2021), *sf* v. 1.0.2 (Pebesma 2018), *geobr* v. 1.6.4 (Pereira and Gonçalves 2021), and *ggplot2* v. 3.3.5. (Wickham 2016). The analysis was performed in R v. 4.1.2 (R Core Team 2021).

## Results

### *Proceratophrys sanctaritae* Cruz & Napoli, 2010

Figure 1

**Material examined.** BRAZIL – Bahia • Jequié, Morro do Mara; 13°53'43.46"S, 039°57'21.64"W; 913 m a.s.l.; 29.VI.2018; APA leg.; 1 ♀, MHNJCH1561; • Jequié, Morro do Mara; 13°53'44.09"S, 039°57'20.90"W; 907 m a.s.l.; 30.VIII.2019, APA leg.; 1 ♂ MHNJCH1710; • Jequié, Morro do Mara; 13°53'44.07"S, 039°57'20.81"W; 906 m a.s.l.; 07.XI.2020, DFOB leg.; 1 ♂ MHNJCH1186 • Wenceslau Guimarães, Wenceslau Guimarães State Ecological Station; 13°35'69"S, 039°43'18"W; 700 m a.s.l.; 04.V.2018; MFVN leg.; 1 juvenile, MZUESC22133.

The Morro do Mara individuals were found at night in the highest parts of the mountain, where they were occupying leaf litter 1–20 m from lotic water bodies. The Wenceslau Guimarães individual was captured during the day (16:30 h) and was foraging in the leaf litter in a hillside forest.

**Identification.** *Proceratophrys sanctaritae* (Fig. 1) is a medium-sized frog presenting a broad and elongated zygomatic branch of the squamosal in sutural contact with the maxilla. This character attributes the species to the genus *Proceratophrys* (Cruz and Napoli 2010). We identified our specimens by their body size (adult males SVL 38.4–45.5 mm in *P. sanctaritae*; Cruz and Napoli 2010, and by the presence of a single long, unicuspidate palpebral appendix with a triangular rostral appendix, which distinguishes this species from others of the




genus that potentially may occur in the areas: *P. renalis* (Miranda-Ribeiro, 1920); *P. schirchii* (Miranda-Ribeiro, 1937), and *P. laticeps* Izecksohn & Peixoto, 1981). The SVL of our specimens ranged from 37.62 mm to 39.09 mm.

Our Moro do Mara and Wenceslau Guimarães records of *P. sanctaritae* are 95 km and 56 km, respectively, from the species' type locality (Fig. 2). The species' extent of occurrence is 613 km<sup>2</sup>, formed by a polygon encompassing the three known occurrences; this area does not represent a prediction analysis of the species' distribution or a refined landscape ecology analysis, as ecological niche modeling has not been done. By including altimetry in the map, other locations in Bahia with conditions similar to those known for *P. sanctaritae* can be inferred as potential occurrence areas and providing a starting point for future surveys; these include the mountain chain that extends from Serra do Timbó to Candido Sales, peripheral to the Espinhaço Mountains (Fig. 2).

## Discussion

Brazil sustains the greatest global amphibian diversity (Segalla et al. 2019). However, there is an alarming lack of information concerning the amphibians' occurrence in Bahia, which is highlighted by the frequent publication of notes on geographic distribution of amphibians in the state (Dias et al. 2010, 2011; Camurugi et al. 2010) and the large number of new species described in the last few years (Caramaschi et al. 2013; Dias et al. 2020; Novaes-e-Fagundes et al. 2021). The same pattern is observed throughout all of Brazil, and, according to Guerra (2020), 25% of the anuran species in Brazil have been described in the last 20 years.

The increase in the number of monitored areas in Bahia (Camurugi et al. 2010; Dias et al. 2014; Freitas et al. 2018; Bastos and Zina in press) can reduce the Wallacean deficit of species whose geographic distributions are still restricted to their type localities (Whittaker et al. 2005). For example, *Phasmahyla timbo* Cruz, Napoli



**Figure 2.** *Proceratophrys sanctaritae* distribution map in Bahia, Brazil, with boundaries between the Caatinga and Atlantic Forest biomes. Similar colors represent environments having similar altimetry. MCP (dashed line) = Extent of Occurrence). (Elevation data from SRTM 90 m DEM Digital Elevation Database).



& Fonseca, 2008 was only observed at its type locality at Serra do Timbó until found in 2007 and 2008 by Camurugi et al. (2010) at the Michelin Biological Reserve in Igrapiúna, Bahia. Another example is *P. sanctaritae*, the focus of this study. Known only by its type locality, it is classified in the Red Book of Brazilian Threatened Fauna (ICMBio 2018) as Critically Endangered. The species carries this status as its occurrence was, until now, restricted to approximately 8 km<sup>2</sup> in a severely fragmented region where, as of 2009, only 3% of native vegetation remains, but forest loss has continued in recent years (ICMBio 2018).

The increase of the known geographic distribution of *P. sanctaritae* results in an extent of occurrence less than 5,000 km<sup>2</sup> (Fig. 2), which implies that this species would better fit Endangered. The new area of occurrence is less than 500 km<sup>2</sup>, and the number of locations are less than five (IUCN 2022). Additionally, all sites where the species is known to occurred are under threat by human activities, mainly from severe habitat loss caused by conversion of native forests towards crops and flooding caused by river damming.

Due to the difficult access and unsuitability for agricultural purposes, most forest remnants outside protected areas in central south Bahia are on slopes or mountain tops (Thomas et al. 1998, 2008). Amphibian diversity hotspots in Bahia's Atlantic Forest are generally associated with mountainous locations, such as Serra Bonita (Dias et al. 2014), Serra do Timbó (Freitas et al. 2019a), and Serra da Jibóia (Freitas et al. 2018). These sites present high rates of endemism and are important reservoirs of biological diversity. Anurans should be highlighted among this biological diversity, as they have unique ecological roles and show high rates of endemism in high-altitude environments.

The current distribution of *P. sanctaritae* is a high-altitude Atlantic Forest's remnant corridor in contact with the Caatinga (Fig. 2). Another frog species, *Bokermannohyla capra* Napoli & Pimenta, 2009, has the same restricted distribution, with isolated populations in high-altitude forest fragments. This similarity may be the effect of past events that resulted in a repeated pattern among frog species and even between higher taxa (Santos 2002; Carnaval et al. 2009, 2014; Freitas et al. 2019b; Menezes et al. 2020; Oliveira-Silva et al. 2021). As *P. sanctaritae* is known only from high altitudes, it can be used as a model for studies on the *P. appendiculata* group's phylogeographic history and to tell the history of the Atlantic Forest and northeastern transition areas.

Although we have not done ecological niche modeling, coincidences are seen between *P. sanctaritae* occurrences in altitude patches within the Atlantic Forest biome. The species may occur in other non-protected areas in a continuum of mountains. Unprotected mountainous areas are the sources of rivers and are often the only localities retaining moisture in an Atlantic Forest/Caatinga matrix. The future of the Atlantic Forest and Caatinga, and the species sheltered by these

morphoclimatic domains, will depend on well-planned conservation strategies based on quality information of its remnants and including investigations of its faunal and floral richness and composition. Creation of new protected areas, restoration projects, and management of natural resources will be critical.

## Acknowledgements

We thank Marina Faraulo and Gabriel Brito for help with the fieldwork, and Jilliard and Jafé for hosting our team. We also thank the reviewers and editors for their contributions.

## Authors' Contributions

Conceptualization: JZ, APA. Formal analysis: DFOB, PBAJ. Investigation: DFOB, MFVN, APA. Methodology: MFVN, PBAJ, APA, DFOB. Validation: JZ. Visualization: IRRD. Writing – original draft: APA, JZ. Writing – review and editing: APA, DFOB, JZ, IRRD.

## References

- Bastos DFO, Zina J (2019) Anfíbiofauna nas alturas: levantamento de anfíbios do Morro do Mara, Jequié, Bahia, Brasil. Monography. Universidade Estadual do Sudoeste da Bahia, Jequié, Brazil.
- Bastos DFO, Zina J (in press) Amphibian fauna from an ecotonal and altitude area in the south-central part of the state of Bahia, Brazil. Herpetology Notes.
- Bivand R, Keitt T, Rowlingson B (2021) rgdal: Bindings for the 'Geospatial' Data Abstraction Library. R package version 1.5-21. <https://CRAN.R-project.org/package=rgdal>
- Camurugi F, Lima, TM, Mercês EA, Juncá FA (2010) Anurans of the Reserva Ecológica da Michelin, Municipality of Igrapiúna, state of Bahia, Brazil. Biota Neotropica 10 (2): 305–312. <https://doi.org/10.1590/S1676-06032010000200032>
- Caramaschi U, Orrico VGD, Faivovich, J, Dias IR, Solé M (2013) A new species of *Allophryne* (Anura: Allophrynidae) from the Atlantic Rain Forest biome of eastern Brazil. Herpetologica 69: 480–491. <https://doi.org/10.1655/herpetologica-d-13-00029>
- Carnaval AC, Hickerson MJ, Haddad CFB, Rodrigues MT, Moritz C (2009) Stability predicts genetic diversity in the Brazilian Atlantic Forest hotspot. Science 323 (5915): 785–789. <https://doi.org/10.1126/science.1166955>
- Carnaval AC, Waltari E, Rodrigues MT, Rosauer D, VanDerWal J, Damasceno R, Prates I, Strangas M, Spanos Z, Rivera D, Pie MR, Firkowski CR, Bornschein MR, Ribeiro LF, Moritz C (2014) Prediction of phylogeographic endemism in an environmentally complex biome. Proceedings of the Royal Society B, Biological Sciences 281 (1792): 14–61. <https://doi.org/10.1098/rspb.2014.1461>
- Cruz AGC, Napoli MF (2010) A new species of smooth horned frog, genus *Proceratophrys* Miranda-Ribeiro (Amphibia: Anura: Cycloramphidae), from the Atlantic Rainforest of eastern Bahia, Brazil. Zootaxa 2660 (1): 57–67. <https://doi.org/10.11646/zootaxa.2660.1.5>
- Dias IR, Vilaça TRA, Silva JRS, Barbosa RS, Solé M (2010) Amphibia, Anura, Hylidae, *Trachycephalus nigromaculatus* Tschudi, 1838: distribution extension. Check List 6 (3): 412–413. <https://doi.org/10.15560/6.3.412>
- Dias IR, Medeiros TT, Solé M, Pimenta BVS (2011) Amphibia, Anura, Hylidae, *Bokermannohyla lucianae* (Napoli and Pimenta, 2003): distribution extension and geographic distribution map. Check List 7 (2): 108–109. <https://doi.org/10.15560/7.2.108>



- Dias IR, Medeiros TT, Vila Nova MF, Solé M (2014) Amphibians of Serra Bonita, southern Bahia: a new hotspot within Brazil's Atlantic Forest hotspot. *ZooKeys* 449: 105–130. <https://doi.org/10.3897/zookeys.449.7494>
- Dias IR, Novaes-E-Fagundes G, Neto A, Zina J et al. (2020) A new large canopy-dwelling species of *Phyllodytes* Wagler, 1930 (Anura, Hylidae) from the Atlantic Forest of the state of Bahia, north-eastern Brazil. *PeerJ* 8: e8642. <https://doi.org/10.7717/peerj.8642>
- Dias PHS, Amaro RC, Carvalho-E-Silva AMPT, Rodrigues MT (2013) Two new species of *Proceratophrys* Miranda-Ribeiro, 1920 (Anura; Odontophrynidae) from the Atlantic Forest, with taxonomic remarks on the genus. *Zootaxa* 3682 (2): 277–304. <https://doi.org/10.11646/zootaxa.3682.2.5>
- Freitas MA, Abegg AD, Dias IR, Moraes EPF (2018) Herpetofauna from Serra da Jibóia, an Atlantic Rainforest remnant in the state of Bahia, northeastern Brazil. *Herpetology Notes* 11: 59–72.
- Freitas MA, Abegg AD, Silva TFS, Fonseca PM, Hamdan B, Filadelfo T (2019a) Herpetofauna of Serra do Timbó, an Atlantic Forest remnant in Bahia state, northeastern Brazil. *Herpetology Notes* 12: 245–260.
- Freitas MA, Abegg AD, Araújo DS, Coelho HEA, Azevedo WDS, Chaves MF, Rosa CM, Moura GJB (2019b) Herpetofauna of three “Brejos de Altitude” in the interior of the state of Pernambuco, northeastern Brazil. *Herpetology Notes* 12: 591–602.
- Frost DR (2022) Amphibian species of the world: an online reference. Version 6.1. American Museum of Natural History, New York, USA. <https://doi.org/10.5531/db.vz.0001>
- Guerra V, Jardim L, Llusia D, Marquez R, Bastos RP (2020) Knowledge status and trends in description of amphibian species in Brazil. *Ecological Indicators* 118: 106754. <https://doi.org/10.1016/j.ecolind.2020.106754>
- Hijmans RJ (2021) raster: geographic data analysis and modeling. R package version 3.4-13, 2021. <https://CRAN.R-project.org/package=raster>
- ICMBio (Instituto Chico Mendes De Conservação Da Biodiversidade) (2018) Livro Vermelho da fauna brasileira ameaçada de extinção. Volume V – anfíbios. ICMBio, Brasília, Brazil, 109–111.
- IUCN (International Union for the Conservation of Nature) (2022a) The IUCN Red List of threatened species. Version 2021-3. <https://www.iucnredlist.org>. Accessed on: 02-22-2022.
- IUCN Standards and Petitions Committee (2022b) Guidelines for using the IUCN Red List categories and criteria. Version 15. Standards and Petitions Committee, IUCN, Gland, Switzerland, 14–16. <https://www.iucnredlist.org/resources/redlistguidelines>.
- Menezes L, Batalha-Filho H, Garda AA, Napoli MF (2020) Tiny treefrogs in the Pleistocene: phylogeography of *Dendropsophus oliveirai* in the Atlantic Forest and associated enclaves in northeastern Brazil. *Journal of Zoological Systematics and Evolutionary Research*. 59 (1): 179–194. <https://doi.org/10.1111/jzs.12422>
- Novaes-E-Fagundes GN, Araujo-Vieira K, Entiauspe-Neto O, Roberto IJ, Orrico VGD, Solé M, Haddad CFB, Loebmann D (2021) A new species of *Scinax* Wagler (Hylidae: Scinaxini) from the tropical forests of northeastern Brazil. *Zootaxa* 4903 (1): 1–41. <https://doi.org/10.11646/zootaxa.4903.1.1>
- Oliveira-Silva W, Siqueira-De-Paula A, Lima HF, Carvalho MGJ, Gaban-Lima R, Lyra-Neves RM, Telino-Junior WR (2021) Avifauna em dois remanescentes de brejo de altitude no estado de Pernambuco, Brasil. *Oecologia Australis* 26 (1): 45–51. <https://doi.org/10.4257/oeco.2022.2601.05>
- Olson DM, Dinerstein E, Wikramanayake ED, Burgess ND, George V. N. Powell, Underwood EC, D'amico JA, Itoua I, Strand HE, Morrison J, Loucks CJ, Allnutt TF, Ricketts T, Kura Y, Lamoreux J, Wettengel WW, Hedao P, Kassem K (2001) Terrestrial ecoregions of the world: a new map of life on Earth. *Bioscience* 51 (11): 933–938. [https://doi.org/10.1641/0006-3568\(2001\)051\[0933:teotwaj\]2.0.co;2](https://doi.org/10.1641/0006-3568(2001)051[0933:teotwaj]2.0.co;2)
- Pebesma E (2018) Simple features for R: standardized support for spatial vector data. *The R Journal* 10 (1): 439–446. <https://doi.org/10.32614/rj-2018-009>
- Pereira RHM, Goncalves CN (2022) geobr: geobr: Download Official Spatial Data Sets of Brazil\_. R package version 1.6.6. <https://github.com/ipeaGIT/geobr>
- Prado GM, Pombal JP (2008) Espécies de *Proceratophrys* Miranda-Ribeiro, 1920 com apêndices palpebrais (Anura; Cycloramphidae). *Arquivos de Zoologia* 39 (1): 1–85. <https://doi.org/10.11606/issn.2176-7793.v39i1p1-85>
- R Core Team (2021) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
- Santana DJ, Silva LAD, Sant'anna AC, Shepard DB, Mângia S (2021) A new species of *Proceratophrys* Miranda-Ribeiro, 1920 (Anura, Odontophrynidae) from southern Amazonia, Brazil. *PeerJ* 9: e12012. <https://doi.org/10.7717/peerj.12012>
- Santos AMM (2002) Distribuição de plantas lenhosas e relações históricas entre a floresta Amazônica, a floresta Atlântica costeira e os brejos de altitude do nordeste brasileiro. Master's dissertation, Universidade Federal de Pernambuco, Recife, Brazil, 77 pp.
- Segalla MV, Caramaschi U, Cruz CAG, Garcia PCA, Grant T, Haddad CFB, Santana DJ, Toledo LF, Langone JA et al. (2019) List of Brazilian amphibians. *Herpetologia Brasileira* 10 (1): 121–216. <https://doi.org/10.5281/zenodo.4716176>
- Thomas WMW, Carvalho AMV, Amorim AMA, Garrison J, Albeláez AL (1998) Plant endemism in two forests in southern Bahia, Brazil. *Biodiversity and Conservation* 7: 311–322. <https://doi.org/10.1023/A:1008825627656>
- Thomas WMW, Carvalho AMV, Amorim AMA, Hanks JG, Santos TS (2008) Diversity of woody plants in the Atlantic coastal forest of southern Bahia, Brazil. *Memoirs of the New York Botanical Garden* 100: 21–66.
- Whittaker RJ, Araújo MB, Jepson P, Ladle RJ, Watson JEM, Willis KJ (2005) Conservation biogeography: assessment and prospect. *Diversity and Distributions* 11: 3–23. <https://doi.org/10.1111/j.1366-9516.2005.00143.x>
- Wickham H (2016) ggplot2: elegant graphics for data analysis. Springer-Verlag, New York, USA. <https://ggplot2.tidyverse.org>